

IN THE CLAIMS

The following is a listing of the claims in the application with claims 1 shown as currently amended, claim 2 shown as deleted and claim 4 shown as withdrawn.

LISTING OF CLAIMS:


1. (currently amended) The fungicidal composition, which comprises:

(i) ~~an agrochemical selected from the group consisting of methyl (2E)-3-methoxy-2-[2'-[[[3''-(1'''-fluoro-2'''-phenyl-1'''-ethenyloxy)phenyl]methylimino]oxy]methylphenyl]propenoate of Formula I); N-methyl(2E)-2-methoxyimino-2-[2'-[[[3''-(1'''-fluoro-2'''-phenyl-1'''-ethenyloxy)phenyl]methylimino]oxy]methylphenylacetamide of Formula (II); and a mixture thereof, and~~


(ii) ~~an adjuvant one or more adjuvant(s) for enhancing the fungicidal efficacy, selected from the group consisting of: a nonionic surfactant selected from the group consisting of a polyoxyethylene-based nonionic surfactant consisting of: a lipophilic moiety being an aliphatic alcohol, a fatty acid or a triacyl glyceride, containing at least 8 carbon atoms, and a hydrophilic moiety being a polyoxyethylene having 3 to 25 oxyethylene repeating units; ethoxylated castor oil; polyoxyethylene-polyoxypropylene block copolymer; and a mixture thereof; an anionic surfactant selected from the group consisting of sodium dioctyl sulfosuccinate, sodium dodecylbenzenesulfonate and a mixture thereof; and a fatty acid alkyl ester;~~

polyoxyethylene lauryl ether, polyoxyethylene cetyl ether, polyoxyethylene stearyl ether and polyoxyethylene oleyl ether, which have 3 to 25 oxyethylene repeating units,

wherein said agrochemical and said adjuvant for enhancing the fungicidal efficacy being mixed in a ratio range from 1:0.5 to 1:20 by weight:



(I)



2. (deleted)
3. (original) The composition of claim 1, which further comprises another agrochemical for preventing or treating plant diseases.
4. (Withdrawn) An adjuvant composition for enhancing the fungicidal efficacy of an agrochemical of compound of Formula (I) or (II) against plant diseases, which comprises:
 - (a) an effective amount of one or more adjuvant for enhancing the fungicidal efficacy selected from the group consisting of a polyoxyethylene-based nonionic surfactants which has an aliphatic alcohol, a fatty acid or triacyl glyceride as a lipophilic moiety containing at least 8 carbon atoms and a polyoxyethylene as a hydrophilic moiety having 3 to 25 oxyethylene repeating units; a polyoxyethylene-polyoxypropylene block copolymer containing 2 to 40 oxyethylene and 25 to 45 oxypropylene repeating units and a mixture thereof; an anionic surfactant selected from the group consisting of sodium dioctyl sulfosuccinate, sodium dodecylbenzenesulfonate and a mixture thereof; and fatty acid alkyl esters having at least 14 carbon atoms;
 - (b) an emulsifier or dispersant; and
 - (c) a carrier.
5. (previously amended) A method for using a fungicidal composition as defined in claims 1 or 2 for preventing or treating a plant disease which comprises spraying the composition at a concentration of 4 to 400 mg/l of the agrochemical or 50 to 2,000 mg/l of the adjuvant for enhancing the fungicidal efficacy to said plant.